

A new pathway of preparation and refined structure of $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$

Darina Lacková, Iveta Ondrejkočová and Marián Koman

Institute of Inorganic Chemistry, Technology and Materials, Faculty of Chemical and Food Technology, Slovak University of Technology, Radlinského 9, 812 37 Bratislava, Slovakia
iveta.ondrejkočovicova@stuba.sk

Abstract: The title compound, $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$ has been prepared by reaction between iron(III) chloride and ammonium chloride which was formed by hydrolysis of isonicotinamide or thionicotinamide in ethanol. The characterization was based on elemental analysis and infrared spectra. The crystal structure of the title compound has been refined by single crystal X-ray diffraction method at 293 K. Crystals are orthorhombic, *Pnma*, with unit cell parameters: $a = 13.760(1) \text{ \AA}$, $b = 9.960(1) \text{ \AA}$, $c = 7.060(1) \text{ \AA}$, $Z = 4$ and $R = 3.5 \%$. The iron(III) atom in $[\text{FeCl}_5(\text{H}_2\text{O})]^{2-}$ ion is approximately octahedrally coordinated by five Cl atoms and one O atom of H_2O molecule. This distortion is caused by the extensively hydrogen-bonded lattice. Structural parameters and IR spectra of similar compounds are compared and discussed.

Keywords: ammonium salt, aqua-pentachloroferrate(III), crystal structure, iron, IR-spectrum

Introduction

The title compound is a member of the series of iron(III) compounds of the general formula $\text{A}_2[\text{FeX}_5(\text{H}_2\text{O})]$, where A is an alkali metal, ammonium or oxonium ion, and X is halide. Their crystallographic and magnetic properties were previously reviewed (Carlin and Palacio 1985). These compounds have been easily prepared from aqueous solutions containing ACl or ABr and appropriate ferric halide, and in several cases also HX . Another compounds which contain $[\text{FeCl}_5(\text{H}_2\text{O})]^{2-}$ anions and organic cations such as are $\text{H}_4(\text{cyclam})^{4+}$, $(\text{enH}_2)^{2+}$ and $(\text{dienH}_3)^{3+}$ have been recently prepared and characterized (Subramanian 1993, James 2001, 2007).

The crystals of $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$ have been usually prepared by slow evaporation of aqueous solutions containing a (1–2):1 molar mix of NH_4Cl and $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (Sharma 1974, Figgis 1978, McElearney 1978, Partiti 1985). This compound was also obtained from hydrochloric acid solutions (Attenborough 1997).

This paper is a continuation of our previously reported studies on spectral and structural properties of iron complexes with derivatives of pyridine (Ondrejkočová et al. 2008, 2009; Štefániková et al. 2008). Here we describe a new pathway of preparation of the title compound and its structural data are compared with published ones.

Experimental

Chemicals and synthesis

All chemicals were purchased commercially and used without further purification. 96 % vol. ethanol was used as a solvent.

The title compound was prepared according following procedure. To an ethanol solution (30 ml) of isonicotinamide, *inia* (1.84 g, 0.015 mol) was added conc. HCl (1.3 ml, 0.015 mol) and a solution which was prepared from $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (4.2 g, 0.015 mol) and ethanol (135 ml). The reaction mixture was stirred for 15 minutes and then left to stand at room temperature. After 3 days, a small amount of precipitated $[\text{iniaH}]\text{Cl}$ was filtered out. Orange crystals of the title compound were obtained from the light orange filtrate after several weeks.

Anal. Calcd. for $[\text{iniaH}]\text{Cl}$ (%): C, 45.44; H, 4.45; N, 17.66. Found: C, 45.70; H, 4.64; N, 17.95. Anal. Calcd. for $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$ (%): H, 3.48; N, 9.74. Found: H, 3.52; N, 9.44.

The title compound was also obtained when thionicotinamide (*tnia*) instead of *inia* was used. There $[\text{tniaH}]\text{Cl}$ was isolated as a by product. Anal. Calcd. for $[\text{tniaH}]\text{Cl}$ (%): C, 41.26; H, 4.04; N, 16.04. Found: C, 40.76; H, 4.06; N, 15.72.

X-ray Crystallography


Intensity data for the title compound was collected using a Siemens P4 diffractometer with graphite monochromated $\text{MoK}\alpha$ radiation (Siemens 1990). The diffraction intensities were corrected for Lorentz and polarization effects with XSCANS (Siemens 1994). Absorption correction was applied using the program XEMP (Sheldrick 1990). The structure was solved by direct methods using the program SHELXS-86 (Sheldrick 1985) and refined by the full-matrix least-squares method on all F^2 data using the program SHELXL-97 (Sheldrick 2008). Geometrical analysis was performed using SHELXL-97.

Tab. 2. Selected bond lengths (Å) and angles (°) for $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$.

Fe—Cl(1)	2.396(2)	Cl(1)—Fe—Cl(2)	92.22(6)	Cl(2)—Fe—O	180.0(2)
Fe—Cl(2)	2.333(2)	Cl(1)—Fe—Cl(3)	179.24(7)	Cl(3)—Fe—Cl(4)	90.49(4)
Fe—Cl(3)	2.363(2)	Cl(1)—Fe—Cl(4)	89.45(4)	Cl(3)—Fe—O	89.4(2)
Fe—Cl(4)	2.390(1)	Cl(1)—Fe—O	89.8(2)	Cl(4)—Fe—O	85.59(4)
Fe—Cl(4A)	2.390(1)	Cl(2)—Fe—Cl(3)	90.54(6)	Cl(4)—Fe—Cl(4A)	171.12
Fe—O	2.119(5)	Cl(2)—Fe—Cl(4)	94.41(3)	H(1)—O—H(1A)	89.29

Tab. 3. Hydrogen bonds with $\text{H}\cdots\text{A} < r(\text{A}) + 2.000 \text{ \AA}$ and $\langle \text{D—H}\cdots\text{A} \rangle > 110^\circ$ for $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$.

D—H	d(D—H)	d(H...A)	$\langle \text{D—H}\cdots\text{A} \rangle$	d(D...A)	A
O1—H1	0.806	2.424	162.30	3.200	Cl4 $[-x, -y, -z + 1]$
N1—H11	0.981	2.521	153.55	3.427	Cl2 $[-x + 1/2, -y, z + 1/2]$
N1—H11	0.981	2.790	114.30	3.316	Cl3 $[-x + 1/2, -y, z - 1/2]$
N1—H12	0.919	2.776	167.01	3.678	Cl4 $[-x + 1/2, -y, z - 1/2]$
N1—H13	0.755	2.780	168.99	3.524	Cl3
N1—H14	0.891	2.833	125.71	3.430	Cl1 $[-x + 1/2, -y, z + 1/2]$
N1—H14	0.891	2.907	133.51	3.579	Cl4 $[x+1/2, y, -z+1/2]$

**Fig. 2.** Packing of molecules in crystal structure of $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$.

The refinement structure of $(\text{NH}_4)_2[\text{FeCl}_5(\text{H}_2\text{O})]$ confirmed the general features published earlier (Figgis 1978). Crystal data and conditions of data collection and refinement are reported in Table 1, bonds lengths and bonds angles are in Table 2 and hydrogen bonds parameters are in Table 3.

The compound is built from one $[\text{FeCl}_5(\text{H}_2\text{O})]^{2-}$ and from two NH_4^+ ions, which are held together by an ionic interaction and additional hydrogen bond inter-

actions between coordinated Cl ligands and ammonia cations, $\text{Cl}\cdots\text{H—N}$. Hydrogen bonds $\text{Cl}\cdots\text{H—O}$ occur also between H_2O and Cl ligands of neighbouring anions (Fig. 1 and Table 3). The system of hydrogen bonds is illustrated in Fig. 1. Packing of the molecules in crystal structure is shown in Fig. 2.

The iron(III) atom in $[\text{FeCl}_5(\text{H}_2\text{O})]^{2-}$ ion is in an approximately octahedral ligand environment formed by five Cl atoms and one molecule of water. This

Tab. 4. Comparison of IR spectra data for $A_2[FeCl_5(H_2O)]$ ($A^+ = NH_4^+, K^+, Rb^+, Cs^+$).

Vibration (cm^{-1})	$(NH_4)_2[FeCl_5(H_2O)]^a$	$K_2[FeCl_5(H_2O)]^b$	$Rb_2[FeCl_5(H_2O)]^b$	$Cs_2[FeCl_5(H_2O)]^b$
$\gamma(NH_4^+)$	1390	-	-	-
$\nu(NH_4^+)$	3184	-	-	-
$\sigma(HOH)$	1582	1585	1584	1593
$\nu_s(OH)$	3388	3369	3391	3389
$\sigma(Fe-Cl)$	279	285, 271	285, 270	not given
$\nu(Fe-O)$	436, 349	376	371	not given

^athis work; ^bPiszczek 2003; Nakamoto 2009.

distortion is caused by the extensively hydrogen-bonded lattice. The Fe—Cl bond (2.333(2) Å) which is *trans* to the water molecule, is evidently shorter than other four Fe—Cl distances (their average is 2.385 Å), presumably as a manifestation of *trans* influence (Table 2). The angles in the anion are very close to the published data (Figgis 1978).

This study is in a good agreement with the earlier X-ray crystallographic reports for $A_2[FeCl_5(H_2O)]$, where A^+ is NH_4^+ , K^+ , Cs^+ , Rb^+ and H_3O^+ (Figgis 1978, McElearney 1978, O'Connor 1979, Søtøfte 1981). When A^+ cation is replaced with the organic cation such as $(H_4(cyclam))^{4+}$, $(enH_2)^{2+}$ or $(dienH_3)^{3+}$, some quite remarkable perturbations from the octahedron were observed (Subramanian 1993, James 2007).

Comparison of the important infrared spectral data for $A_2[FeCl_5(H_2O)]$, where A^+ is NH_4^+ , K^+ , Rb^+ or Cs^+ are reported in Table 4. The most intense bands in the spectrum are at 3184 and 1390 cm^{-1} . These bands can be assigned to NH_4^+ vibrations (Nakamoto 2009). The $\delta(HOH)$ and $\nu_s(OH)$ bands which occur at 1582 and 3388 cm^{-1} , respectively clearly confirm the presence of water molecule in the anion (Piszczek 2003). Other bands at 436 and 349 cm^{-1} confirm the bond formation between iron and oxygen atoms. The peak at 279 cm^{-1} is attributed to the $\delta(Fe-Cl)$, what is also in a good agreement with the literature data for the similar compounds (Piszczek 2003; Nakamoto 2009).

Acknowledgement

This work was supported by courtesy of the Slovak Grant Agency (VEGA 1/0056/13 and APVV-0014-11).

References

Attenborough M, Hall I, Nikolov O, Brown SR, (1997) *Hyperfine Interactions* 108: 435–442.
Carlin RL, Palacio F (1985) *Coord. Chem. Rev.* 65: 141–165.
Farrugia LJ (1997) *J. Appl. Cryst* 30: 565–566.
Farrugia LJ (1999) *J. Appl. Cryst* 32: 837–838.

Figgis BN, Raston CL, Sharma RP, White AH (1978) *Aust. J. Chem.* 31: 2717–2720.
James BD, Bakalova M, Liesegang J, Feiff WM, Skelton BW, White AH (2001) *Inorg. Chem.* 40: 4617–4622.
James BD, Mrozinski J, Klak J, Skelton BW, White AH, (2007) *Z. Anorg. Allg. Chem.* 663: 2683–2688.
James BD, Mrozinski J, Klak J, Skelton BW, White AH (2007) *Z. Anorg. Allg. Chem.* 633: 974–978.
Jellinek HHG, Urwin JR (1953) *J. Phys. Chem.* 57: 900–902.
Macrae CF, Edgington PR, McCabe P, Pidcock E, Shields GP, Taylor R, Towler M, van de Streek J (2006) *J. Appl. Cryst* 39: 453–457.
McElearney JN, Merchant S (1978) *Inorg. Chem.* 17: 1207–1215.
Nakamoto Kazuo (2009). In: *Infrared and Raman Spectra of Inorganic and Coordination Compounds; Part A: Theory and Applications in Inorganic Chemistry*, Wiley.
O'Connor ChJ, Deaver BS Jr, Sinn E (1979) *J. Chem. Phys.* 70: 5161–5167.
Ondrejčková I, Galková S, Mrozinski J, Klak J, Lis T, Olejnik Z (2008) *Inorg. Chim. Acta* 361: 2483–2490.
Ondrejčková I, Wrzeczion M, Nahorska M, Mrozinski J (2009) *Polish J. Chem.* 83: 1547–1553.
Partiti CSM, Piccini A, Rechenberg HR (1985) *Solid State Commun.* 56: 687–691.
Piszczek P, Grodzicki A, Engelen B (2003) *J. Mol. Struct.* 646: 45–54.
Sharma SK, Pandya DK (1974) *J. Inorg. Nucl. Chem.* 36: 1165–1166.
Sheldrick GM (1990) *Acta Cryst.* A46: 467–473.
Sheldrick GM (1985) SHELXS-86. In: Sheldrick, GM, Krueger R, Goddard R (Eds) *Crystallographic Computing 3*, Oxford University Press, 175.
Sheldrick GM (2008) *Acta Cryst.* A64: 112–122.
Siemens (1990) XEMP. Version 4.2, Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
Siemens (1994) Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
Smith BC, Anderson MA, Hoadley KA, Keck JL, Cleland WW, Denu JM (2012) *Biochemistry* 51: 243–256.
Søtøfte I, Nielsen K (1981) *Acta Chem. Scand. A* 35: 821–822.
Subramanian S, Zaworotko MJ (1993) *Can. J. Chem.* 71: 433–440.
Štefánková S, Ondrejčková I, Koman M, Lis T, Mrozinski J, Wrzeczion (2008) *J. Coord. Chem.* 61: 3895–3903.